

Alavert Allergy Citrus 60ct Tablets

Drug Facts

Active ingredient (in each tablet)	Purpose
Loratadine 10 mg.....	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing
■ itchy, watery eyes ■ itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not use more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ Place 1 tablet on tongue; tablet disintegrates with or without water.

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- phenylketonurics: contains phenylalanine 0.56 mg per tablet
- safety sealed: do not use if the individual blister unit imprinted with Alavert® Citrus Burst is open or torn
- store between 20-25°C (68 to 77°F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients anhydrous citric acid, aspartame, flavors, mannitol, polysorbate 80, pullulan

Questions or comments?

Call 1-800 ALAVERT (1-800-252-8378) weekdays 9 AM to 5 PM EST