

Alavert D-12 Hour 24ct Tablets

<p>Drug Facts</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black;">Active ingredients (in each tablet)</td> <td style="width: 50%; border-bottom: 1px solid black;">Purposes</td> </tr> <tr> <td>Loratadine 5 mg.....</td> <td>Antihistamine</td> </tr> <tr> <td>Pseudoephedrine sulfate 120 mg.....</td> <td>Nasal decongestant</td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: <ul style="list-style-type: none"> ■ sneezing ■ itchy, watery eyes ■ runny nose ■ itching of the nose or throat ■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ reduces swelling of nasal passages ■ temporarily relieves sinus congestion and pressure ■ temporarily restores freer breathing through the nose <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ liver or kidney disease. Your doctor should determine if you need a different dose. 	Active ingredients (in each tablet)	Purposes	Loratadine 5 mg.....	Antihistamine	Pseudoephedrine sulfate 120 mg.....	Nasal decongestant	<p>Drug Facts (continued)</p> <p>When using this product do not take more than directed. Taking more than directed may cause drowsiness.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve within 7 days or are accompanied by a fever ■ nervousness, dizziness or sleeplessness occurs <p>if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not divide, crush, chew or dissolve the tablet <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;">adults and children</td> <td style="width: 50%;">1 tablet every 12 hours;</td> </tr> <tr> <td>12 years and over</td> <td>not more than 2 tablets in 24 hours</td> </tr> <tr> <td>children under 12 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> ■ each tablet contains: calcium 25 mg ■ store between 20° to 25°C (68° to 77°F) ■ keep in a dry place <p>Inactive ingredients croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide</p> <p>Questions or comments? Call 1-800 ALAVERT (1-800-252-8378) weekdays 9 AM to 5 PM EST</p>	adults and children	1 tablet every 12 hours;	12 years and over	not more than 2 tablets in 24 hours	children under 12 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor
Active ingredients (in each tablet)	Purposes														
Loratadine 5 mg.....	Antihistamine														
Pseudoephedrine sulfate 120 mg.....	Nasal decongestant														
adults and children	1 tablet every 12 hours;														
12 years and over	not more than 2 tablets in 24 hours														
children under 12 years of age	ask a doctor														
consumers with liver or kidney disease	ask a doctor														